

On globally nilpotent differential equations

Michael Dettweiler and Stefan Reiter

February 2, 2008

Abstract

In a previous work of the authors, a middle convolution operation on the category of Fuchsian differential systems was introduced. In this note we show that the middle convolution of Fuchsian systems preserves the property of global nilpotence.

This leads to a globally nilpotent Fuchsian system of rank two which does not belong to the known classes of globally nilpotent rank two systems. Moreover, we give a globally nilpotent Fuchsian system of rank seven whose differential Galois group is isomorphic to the exceptional simple algebraic group of type G_2 .

Introduction

A unifying description of all irreducible and physically rigid local systems on the punctured affine line was given by Katz [9]. The main tool therefore is a middle convolution functor on the category of perverse sheaves (loc. cit., Chap. 5). In [4], the authors give a purely algebraic analogon of this convolution functor. This functor is a functor of the category of finite dimensional K -modules of the free group F_r on r generators to itself (K denoting a field). It depends on a scalar $\lambda \in K^\times$ and is denoted by MC_λ .

By the Riemann Hilbert correspondence (see [2]), a construction parallel to MC_λ should exist in the category of Fuchsian systems of differential equations. In [5], such a construction is given, leading to a description of rigid Fuchsian systems which is parallel to Katz description of rigid local systems. The convolution depends on a parameter $\mu \in \mathbb{C}$ and carries a Fuchsian system F to another Fuchsian system, denoted by $\mathrm{mc}_\mu(F)$, see Section 1.2.

In this note we study how the $\mathfrak{p}$ -curvature (for $\mathfrak{p}$ a prime of a number field K) of a Fuchsian system F having coefficients in the function field $K(t)$ changes under

the convolution process. The $\mathfrak{p}$ -curvature is a matrix $\bar{C}_{\mathfrak{p}}(F)$ with coefficients in the function field over a finite field which is obtained from a p -fold iteration of F (where p is the prime number below $\mathfrak{p}$) and reduction modulo $\mathfrak{p}$, see Section 2. The $\mathfrak{p}$ -curvature matrices encode many arithmetic and geometric properties of a Fuchsian system. For example, the Bombieri-Dwork conjecture predicts that if the $\mathfrak{p}$ -curvature $\bar{C}_{\mathfrak{p}}(F)$ is nilpotent for almost all primes $\mathfrak{p}$ of K (i.e., F is *globally nilpotent*), then F is arising from geometry, see [1]. In this note we prove the following result (see Thm. 2.0.6):

Theorem 1: *Let F be a Fuchsian system, let $\mu \in \mathbb{Q}$ and let $\mathrm{mc}_{\mu}(F)$ be the middle convolution of F with respect to μ . Then the following holds:*

- (i) *If $\mathrm{ord}_p(\mu) \geq 0$ and if the p -curvature of F is nilpotent of rank k , then the p -curvature of the middle convolution $\mathrm{mc}_{\mu}(F)$ is nilpotent of rank $r \in \{k-1, k, k+1\}$.*
- (ii) *If F is globally nilpotent, then $\mathrm{mc}_{\mu}(F)$ is globally nilpotent.*

Of course, the second statement of the theorem follows immediately from the first. The second statement can be deduced alternatively from the stability of global nilpotence under pullback, tensor product, higher direct image, see Katz [8], Section 5.7 through 5.10 (it follows from [5], Thm. 1.2, and [3], Rem. 3.3.7, that mc_{μ} corresponds under the Riemann-Hilbert correspondence to the middle convolution MC_{χ} of local systems $\mathcal{V}$ with Kummer sheaves - which is, by construction, a higher direct image sheaf).

The proof of the first statement relies on the closed formula of the p -curvature of Okubo systems (compare to Remark 2.0.4) and on the fact that the middle convolution of a Fuchsian system is a factor system of an Okubo system.

In the last section, we apply the above result in order to obtain two new globally nilpotent differential equations:

In Section 3.1, we apply the middle convolution to the globally nilpotent Fuchsian system of rank two which appears in the work of Krammer [10]. This leads again to a globally nilpotent Fuchsian system of rank two. It is a new type of a globally nilpotent rank two system because it is neither a pullback of a hypergeometric system, nor a system associated to periods of Shimura curves (see Thm. 3.1.1 and the discussion in [10], Section 11).

In Section 3.2 we consider a globally nilpotent Fuchsian system G of rank seven which is constructed by an sixfold iteration of middle convolutions and tensor products. Using the Riemann-Hilbert correspondence for G (see Thm. 1.2.3), one

can see that the Zariski closure of monodromy group of G is equal to the simple exceptional algebraic group of type G_2 . The Fuchsian system G can be viewed as the de Rham version of recent results of the authors on rigid local systems and motives with Galois group G_2 ([6]). The motivic interpretation of the rigid local system which is given there, and the Riemann-Hilbert correspondence (see Thm. 1.2.3), imply that the Bombieri-Dwork conjecture holds for the Fuchsian system G .

The authors thank J. Aidan for pointing out a misprint in the differential equation which appears in Thm. 3.1.1 (in an earlier version of this paper) and N. Katz for valuable comments on the nilpotence of higher direct image connections.

1 The Riemann-Hilbert correspondence of the middle convolution

1.1 The tuple transformation MC_λ . Let K be a field, let V be a finite dimensional vectorspace over K and let $M = (M_1, \dots, M_r)$ be an element of $GL(V)^r$. For any $\lambda \in K^\times$ one can construct another tuple of matrices $(N_1, \dots, N_r) \in GL(V^r)^r$, as follows: For $k = 1, \dots, r$, N_k maps a vector $(v_1, \dots, v_r)^{\text{tr}} \in V^r$ to

$$\begin{pmatrix} 1 & 0 & & \dots & & 0 \\ & \ddots & & & & \\ & & 1 & & & \\ (M_1 - 1) & \dots & (M_{k-1} - 1) & \lambda M_k & \lambda(M_{k+1} - 1) & \dots & \lambda(M_r - 1) \\ & & & 1 & & & \\ & & & & \ddots & & \\ 0 & & & \dots & & 0 & 1 \end{pmatrix} \begin{pmatrix} v_1 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ v_r \end{pmatrix}.$$

There are the following $\langle N_1, \dots, N_r \rangle$ -invariant subspaces of V^r :

$$\mathcal{K}_k = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \ker(M_k - 1) \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad (k\text{-th entry}), \quad k = 1, \dots, r,$$

and

$$\mathcal{L} = \cap_{k=1}^r \ker(N_k - 1) = \ker(N_r \cdots N_1 - 1).$$

Let $\mathcal{K} := \oplus_{i=1}^r \mathcal{K}_i$.

1.1.1 Definition. Let $\text{MC}_\lambda(M) := (\tilde{N}_1, \dots, \tilde{N}_r) \in \text{GL}(V^r/(\mathcal{K} + \mathcal{L}))^r$, where $\tilde{N}_k$ is induced by the action of N_k on $V^r/(\mathcal{K} + \mathcal{L})$. We call $\text{MC}_\lambda(M)$ the *middle convolution* of M with λ .

1.2 The middle convolution of Fuchsian systems Let $A = (A_1, \dots, A_r)$, $A_k \in \mathbb{C}^{n \times n}$. For $\mu \in \mathbb{C}$ one can define blockmatrices B_k , $k = 1, \dots, r$, as follows:

$$B_k := \begin{pmatrix} 0 & & \dots & 0 \\ & \ddots & & \\ A_1 & \dots & A_{k-1} & A_k + \mu & A_{k+1} & \dots & A_r \\ & & & \ddots & & & \\ 0 & & \dots & & & & 0 \end{pmatrix} \in \mathbb{C}^{nr \times nr},$$

where B_k is zero outside the k -th block row.

The tuple

$$(1.2.1) \quad c_\mu(A) := (B_1, \dots, B_r)$$

is called the *naive convolution* of A with μ . There are the following left- $\langle B_1, \dots, B_r \rangle$ -invariant subspaces of the column vector space $\mathbb{C}^{nr}$ (with the tautological action

of $\langle B_1, \dots, B_r \rangle$:

$$\mathfrak{k}_k = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \ker(A_k) \\ 0 \\ \vdots \\ 0 \end{pmatrix} \quad (k\text{-th entry}), \quad k = 1, \dots, r,$$

and

$$\mathfrak{l} = \cap_{k=1}^r \ker(B_k) = \ker(B_1 + \dots + B_r).$$

Let $\mathfrak{k} := \oplus_{k=1}^r \mathfrak{k}_k$ and fix an isomorphism between $\mathbb{C}^{nr}/(\mathfrak{k} + \mathfrak{l})$ and $\mathbb{C}^m$.

1.2.1 Definition. The tuple of matrices $\text{mc}_\mu(A) := (\tilde{B}_1, \dots, \tilde{B}_r) \in \mathbb{C}^{m \times m}$, where $\tilde{B}_i$ is induced by the action of B_i on $\mathbb{C}^m (\simeq \mathbb{C}^{nr}/(\mathfrak{k} + \mathfrak{l}))$, is called the *middle convolution* of A with μ .

Let $S := \mathbb{C} \setminus \{t_1, \dots, t_r\}$ and $A := (A_1, \dots, A_r)$, $A_i \in \mathbb{C}^{n \times n}$. The Fuchsian system

$$F : Y' = \sum_{i=1}^r \frac{A_i}{t - t_i} Y$$

is called *the Fuchsian system associated to the tuple A* .

1.2.2 Definition. Let $A := (A_1, \dots, A_r)$, $A_i \in \mathbb{C}^{n \times n}$, and $\mu \in \mathbb{C}$. Let F be the Fuchsian system associated to A . Then the Fuchsian system F which is associated to the middle convolution tuple $\text{mc}_\mu(A)$ is denoted by $\text{mc}_\mu(F)$ and is called the *middle convolution* of F with the parameter μ . The Fuchsian system which is associated to the naive convolution tuple $\text{c}_\mu(A)$ is denoted by $\text{c}_\mu(F)$ and is called the *naive convolution* of F with the parameter μ .

Fix a set of homotopy generators δ_i , $i = 1, \dots, r$, of $\pi_1(S, t_0)$ by traveling from the base point t_0 to t_i , then encircling t_i counterclockwise, and then going back to t_0 . Then, the analytic continuation of solutions along δ_i defines a linear isomorphism M_i , $i = 1, \dots, r$, of the vectorspace $V \simeq \mathbb{C}^n$ of holomorphic solutions of F which are defined in a small neighborhood of t_0 . We call the tuple $(M_1, \dots, M_r)$

the *monodromy tuple* of F . The Riemann-Hilbert correspondence says that F is determined up to isomorphism by M , see [2].

The following result is an explicit realization of the Riemann-Hilbert correspondence for a convoluted Fuchsian system, see [5], Thm. 6.8:

1.2.3 Theorem. *Let F be an irreducible Fuchsian system associated to $A = (A_1, \dots, A_r) \in (\mathbb{C}^{n \times n})^r$. Assume that there exist two different elements such that the local monodromy of $\mathcal{V}$ is nontrivial. Fix a set of homotopy generators*

$$\delta_1, \dots, \delta_r \in \pi_1(\mathbb{A}^1 \setminus \{t_1, \dots, t_r\})$$

of the fundamental group as above. Let $M = (M_1, \dots, M_r) \in \mathrm{GL}_n(\mathbb{C})^r$ be the monodromy tuple of F (with respect to $\delta_1, \dots, \delta_r$). Assume that

$$\mathrm{rk}(A_i) = \mathrm{rk}(M_i - 1), \quad \mathrm{rk}(A_1 + \dots + A_r + \mu) = \mathrm{rk}(\lambda \cdot M_1 \cdots M_r - 1).$$

Then the monodromy tuple of $\mathrm{mc}_\mu(F)$ is given by $\mathrm{MC}_\lambda(M)$, where $\lambda = e^{2\pi i \mu}$.

2 Transformation of the p -curvature under mc_μ

Let K be a number field and let $F : Y' = CY$ be a system of linear differential equations, where $C \in K(t)^{n \times n}$. Successive application of differentiation yields differential systems

$$(2.0.2) \quad F^{(n)} : Y^{(n)} = C_n Y \quad \text{for each } n \in \mathbb{N}_{>0}.$$

In the following, $\mathfrak{p}$ always denotes a prime of K which lies over a prime number p . For almost all primes p , one can reduce the entries of the matrices C_p modulo $\mathfrak{p}$ in order to obtain the $\mathfrak{p}$ -curvature matrices of F

$$\bar{C}_{\mathfrak{p}} = \bar{C}_{\mathfrak{p}}(F) := C_p \pmod{\mathfrak{p}}.$$

2.0.1 Definition. A system of linear differential equations F can be written in *Okubo normal form*, if F can be written as

$$F : Y' = (t - T)^{-1} B Y,$$

where $B \in \mathbb{C}^{n \times n}$ and T is a diagonal matrix $T = \mathrm{diag}(t_1, \dots, t_n)$ with $t_i \in \mathbb{C}$ (here possibly $t_i = t_j$ for $i \neq j$).

The following proposition is obvious from the definitions:

2.0.2 Proposition. *If F is a Fuchsian system, then the naive convolution $c_\mu(F)$ of F can be written in Okubo normal form.*

An induction yields the following formula for the $\mathfrak{p}$ -curvature matrix of a system in Okubo normal form:

2.0.3 Lemma. *Let*

$$F : Y' = CY = (t - T)^{-1}BY$$

be a system of linear differential equations which can be written in Okubo normal form. Then

$$(2.0.3) \quad C_n = (t - T)^{-1}(B - n + 1) \cdot (t - T)^{-1}(B - n + 2) \cdots \\ \cdots (t - T)^{-1}(B - 1) \cdot (t - T)^{-1}B.$$

Epecially, if the matrix B has coefficients in a number field K , then the $\mathfrak{p}$ -curvature $\bar{C}_{\mathfrak{p}}$ of F has the form

$$(t - T)^{-1}(B - p + 1) \cdot (t - T)^{-1}(B - p + 2) \cdots (t - T)^{-1}(B - 1) \cdot (t - T)^{-1}B \pmod{\mathfrak{p}}.$$

□

2.0.4 Remark. The above proposition is interesting because there is no closed formula known for the computation of the $\mathfrak{p}$ -curvature if the rank is > 2 (compare to [13] and [12]). The above lemma yields such a closed formula for Okubo systems. On the other hand, every irreducible Fuchsian system is a subfactor of an Okubo system (this is known by the work of Okubo and follows also from Prop. 2.0.2 and the fact that, by Equation (2.0.4) below, the middle convolution mc_{-1} induces the identity transformation of Fuchsian systems).

The following technical proposition will be used below:

2.0.5 Proposition. *Let K be a number field, let $F : Y' = CY$, $C \in K(t)^{n \times n}$, be a Fuchsian system and let $c_0(F) : Y' = DY$ be the naive convolution of F with the parameter 0. Let p be a prime number and let $\mathfrak{p}$ denote a prime of K which lies over p such that the entries of C can be reduced modulo $\mathfrak{p}$. If $C_p^k \equiv 0 \pmod{\mathfrak{p}}$, then $D_{kp+1} \equiv 0 \pmod{\mathfrak{p}}$.*

Proof: The Fuchsian system F is the Fuchsian system associated to some tuple $(A_1, \dots, A_r) \in (K^{n \times n})^r$. By Prop. 2.0.2, the naive convolution of F with the parameter -1 can be written in Okubo normal form

$$c_{-1}(F) : Y' = (t - T)^{-1}\tilde{B}Y = \tilde{D}Y.$$

Here, T is the diagonal matrix $T = \text{diag}(t_1, \dots, t_1, \dots, t_r, \dots, t_r)$ (every t_k occurs n times) and

$$\tilde{B} = (\tilde{B}_{i,j}), \quad \tilde{B}_{i,j} = A_j - \delta_{i,j} E_n$$

($E_n \in \text{GL}_n(K)$ denoting the identity matrix). Using the base change which is induced by the matrix $H(t - T)$, where

$$H := \begin{pmatrix} E_n & -E_n & 0 & \dots \\ 0 & \ddots & \ddots & \\ \vdots & & & -E_n \\ 0 & \dots & & E_n \end{pmatrix},$$

one can verify that the naive convolution $c_{-1}(F)$ is equivalent to the following system:

$$(2.0.4) \quad Y' = GY = \begin{pmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ \frac{A_1}{t-t_1} & (\frac{A_1}{t-t_1} + \frac{A_2}{t-t_2}) & \dots & (\frac{A_1}{t-t_1} + \dots + \frac{A_r}{t-t_r}) \end{pmatrix} Y.$$

By a straightforward computation, one sees that

$$(2.0.5) \quad G = H(t - T) \cdot D \cdot (H(t - T))^{-1},$$

where D is the matrix appearing in the naive convolution $c_0(F) : Y' = DY$ with the parameter 0. Using the Leibnitz rule, one can further verify that

$$(2.0.6) \quad G_p = (H(t - T)) \cdot \tilde{D}_p \cdot (H(t - T))^{-1}$$

(note that $\tilde{D}_p$ belongs to the naive convolution $c_{-1}(F)$ with parameter -1). It follows from Equation (2.0.4) that

$$G_p = \begin{pmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ * & * & * & C_p \end{pmatrix}.$$

Using this and the assumption $C_p^k \equiv 0 \pmod{\mathfrak{p}}$ one sees that

$$G_p^k G = \begin{pmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ * & * & * & 0 \end{pmatrix} \begin{pmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ * & * & * & * \end{pmatrix} \equiv 0 \pmod{\mathfrak{p}}.$$

Thus, by Equations (2.0.5) and (2.0.6),

$$H(t-T) \cdot \tilde{D}_p^k \cdot D \cdot (H(t-T))^{-1} \equiv 0 \pmod{\mathfrak{p}}$$

and thus

$$\tilde{D}_p^k \cdot D \equiv 0 \pmod{\mathfrak{p}}.$$

The claim follows now from this and the equality

$$\tilde{D}_p^k \cdot D \equiv D_{pk+1} \pmod{\mathfrak{p}},$$

which is immediate from the Equation 2.0.3 appearing in Lemma 2.0.3. $\square$

2.0.6 Theorem. *Let K be a number field, let*

$$F : Y' = CY, \quad C \in K(t)^{n \times n},$$

be a Fuchsian system and let $\mu \in \mathbb{Q}$. Let p be a prime number and let $\mathfrak{p}$ denote a prime of K which lies over p such that the number μ and the entries of C can be reduced modulo $\mathfrak{p}$. If the $\mathfrak{p}$ -curvature of F is nilpotent of rank k (i.e., $\bar{C}_p(F)^k \equiv 0$), then the $\mathfrak{p}$ -curvature of the middle convolution $\text{mc}_\mu(F)$ is nilpotent of rank $r \in \{k-1, k, k+1\}$.

Proof: We first prove that the $\mathfrak{p}$ -curvature of the naive convolution $c_\mu(F)$ is nilpotent of rank at most $k+1$: Let $c_0(F) : Y' = DY$ be the naive convolution of F with the parameter 0 and let $c_\mu(F) : Y' = D^\mu Y$ be the naive convolution of F with the parameter μ . By assumption, one has $C_p^k \equiv 0 \pmod{\mathfrak{p}}$. Thus, by the preceding proposition,

$$(2.0.7) \quad D_{kp+1} \equiv 0 \pmod{\mathfrak{p}}.$$

One can easily deduce from the Equation (2.0.3) of Lemma 2.0.3, that the matrix $D_{kp+1} \pmod{\mathfrak{p}}$ appears as a factor in the product formula (2.0.3) for $D_{p(k+1)}^\mu$ taken modulo $\mathfrak{p}$. Thus, by Equation (2.0.7) and again by Lemma 2.0.3,

$$0 \equiv D_{p(k+1)}^\mu \equiv (D_p^\mu)^{k+1} = \bar{C}_p(\text{mc}_\mu(F))^{k+1} \pmod{\mathfrak{p}}.$$

Since the middle convolution $\text{mc}_\mu(F)$ is a factor of the naive convolution $c_\mu(F)$, it follows that the $\mathfrak{p}$ -curvature of the middle convolution $\text{mc}_\mu(F)$ is nilpotent of rank at most $k + 1$. Now the claim follows from the fact that $\text{mc}_{-\mu} \circ \text{mc}_\mu = \text{id}$. $\square$

3 Two new examples of globally nilpotent Fuchsian systems

3.1 A globally nilpotent Fuchsian system of rank two. Dwork has conjectured that any globally nilpotent second order differential equation has either algebraic solutions or has a correspondence to a Gauß hypergeometric differential equation (see [10]). This conjecture was disproved by Krammer (loc. cit.). The counterexample which Krammer gives is the uniformizing differential equation K of an arithmetic Fuchsian lattice. It comes from the periods of a family of abelian surfaces over a Shimura curve $S \simeq \mathbb{P}^1 \setminus \{0, 1, 81, \infty\}$. The equation K is an irreducible ordinary second order differential equation which can easily be transformed into the following Fuchsian system of rank two:

$$K : Y' = \left(\frac{1}{t} \cdot \begin{pmatrix} 0 & 0 \\ -\frac{1}{2} & -\frac{1}{2} \end{pmatrix} + \frac{1}{t-1} \cdot \begin{pmatrix} 0 & 0 \\ \frac{4}{9} & -\frac{1}{2} \end{pmatrix} + \frac{1}{t-81} \begin{pmatrix} 0 & 1 \\ 0 & \frac{1}{2} \end{pmatrix} \right) \cdot Y.$$

The local monodromy of K at the finite singularities is given by three reflections, the local monodromy of K at ∞ is given by an element of order six.

3.1.1 Theorem. *The middle convolution $H := \text{mc}_{\frac{1}{6}}(K)$ is equivalent to the following Fuchsian system:*

$$Y' = \left(\frac{1}{t} \begin{pmatrix} -\frac{19}{30} & \frac{19}{10} \\ -\frac{1}{10} & \frac{3}{10} \end{pmatrix} + \frac{1}{t-1} \begin{pmatrix} -\frac{1}{3} & -\frac{7}{18} \\ 0 & 0 \end{pmatrix} + \frac{1}{t-81} \begin{pmatrix} 0 & 0 \\ -\frac{1}{2} & \frac{2}{3} \end{pmatrix} \right) Y.$$

The Fuchsian system H is globally nilpotent. Moreover, the system H has neither a correspondence to a hypergeometric differential system, nor a correspondence to a uniformizing differential equation of a Fuchsian lattice.

Proof: The first claim follows from Cor. 2.0.6. To prove the last statement, we use the Riemann-Hilbert correspondence: The monodromy tuple of K can be

shown to be

$$A := \left(i \cdot \begin{pmatrix} 0 & -(\sqrt{5}+1)/2 \\ (\sqrt{5}-1)/2 & 0 \end{pmatrix}, i \cdot \begin{pmatrix} -\sqrt{3} & 1+\sqrt{5} \\ 1-\sqrt{5} & \sqrt{3} \end{pmatrix}, \right. \\ \left. i \cdot \begin{pmatrix} (\sqrt{3}+\sqrt{15})/2 & -2 \\ 2 & (\sqrt{3}-\sqrt{15})/2 \end{pmatrix} \right).$$

In the following, the element ζ_n denotes the root of unity $e^{2\pi i/n}$. Using Thm. 1.2.3, it is easy to see that the monodromy of $\text{mc}_{\frac{1}{6}}$ is given by

$$\text{MC}_{\zeta_6}(A) = (B_1, B_2, B_3) =$$

$$\left(\zeta_6 \cdot \begin{pmatrix} \frac{1}{5}(\sqrt{5} + 3\zeta_{60}^{10} + 6\zeta_{60}^8 - 6\zeta_{60}^2 - 3) & \frac{1}{5}(-6\sqrt{5} - 10\zeta_{60}^{10} - 10\zeta_{60}^8 + 10\zeta_{60}^2 + 10) \\ \frac{1}{5}(\zeta_{60}^{14} + 2\zeta_{60}^{10} - \zeta_{60}^8 - \zeta_{60}^6 - \zeta_{60}^4 + \zeta_{60}^2 + 1) & \frac{1}{5}(-\sqrt{5} - 3\zeta_{60}^{10} - 6\zeta_{60}^8) \end{pmatrix}, \right. \\ \left. \zeta_6 \cdot \begin{pmatrix} -\zeta_{60}^{10} & 2\zeta_{60}^{10} \\ 0 & \zeta_{60}^{10} - 1 \end{pmatrix}, \zeta_6 \cdot \begin{pmatrix} \zeta_{60}^{10} - 1 & 0 \\ -\zeta_{60}^{10} + 1 & -\zeta_{60}^{10} \end{pmatrix} \right).$$

By [4], Cor 5.9, one can show that the elements B_i , $i = 1, 2, 3$, generate a subgroup which is conjugate to a subgroup of $\text{SU}_{1,1}(\mathbb{R})$. The latter group is well known to be conjugate to the group $\text{SL}_2(\mathbb{R})$ inside the group $\text{GL}_2(\mathbb{C})$.

Consider the element

$$\tilde{B} := B_1 B_2 B_1 = \begin{pmatrix} 2\zeta_{60}^{10} - 1 & -4\zeta_{60}^{10} - 2\zeta_{60}^8 + 2\zeta_{60}^2 \\ -\zeta_{60}^{14} + \zeta_{60}^8 + \zeta_{60}^6 + \zeta_{60}^4 - \zeta_{60}^2 - 1 & 2\zeta_{60}^{14} - 2\zeta_{60}^{10} - 2\zeta_{60}^6 - 2\zeta_{60}^4 + 3 \end{pmatrix}.$$

It is an elliptic element since the absolute value of its trace is $|- \sqrt{5} + 1| < 2$. Moreover, the order of $\tilde{B}$ is infinite. (The eigenvalues of $\tilde{B}$ are roots of $f := X^4 - 2X^3 - 2X^2 - 2X + 1$ with $\text{Gal}(f) = D_8$, thus the eigenvalues cannot be roots of unity.) Thus, the monodromy tuple $\text{MC}_{\zeta_6}(A)$ of $\text{mc}_{\frac{1}{6}}(F)$ generates a non-discrete subgroup of $\text{PGL}_2(\mathbb{R})$ and thus does not have any correspondence to a hypergeometric differential system (in which case it would be a discrete triangle group, see [7], 4.1.2), nor a correspondence to a uniformizing differential equation of a Fuchsian lattice. $\square$

3.2 A globally nilpotent system with differential Galois group G_2 .

The following construction of local systems with finite monodromy will be used in Section 3.2 below: Let $\rho_f : \pi_1(X) \rightarrow G$ be the surjective homomorphism associated to a finite Galois cover with Galois group G and let $\alpha : G \rightarrow \mathrm{GL}_n(\mathbb{C})$ be a representation. Then the composition

$$\alpha \circ \rho_f : \pi_1(X) \rightarrow \mathrm{GL}_n(\mathbb{C})$$

corresponds to a local system with finite monodromy on X which is denoted by $\mathcal{V}_{f,\alpha}$.

Consider the following local systems with finite monodromy

$$\mathcal{V}_i = \mathcal{V}_{f_i,\alpha}, \quad i = 0, 1, 2,$$

on $S = \mathbb{A}^1 \setminus \{0, 1\}$, where the f_i and α are as follows: The representation α is the embedding of $\mathbb{Z}/2\mathbb{Z}$ into $\mathbb{Q}_\ell^\times$ and the $f_i : Y_i \rightarrow S$, $i = 0, 1, 2$, are the double covers defined by the equations

$$z^2 = t(t-1), \quad \text{resp.} \quad z^2 = t-1, \quad \text{resp.} \quad z^2 = t.$$

The middle convolution operation we use is MC_χ , where χ is associated to the double cover of $\mathbb{G}_m$ given by $z^2 = t$ and by $\alpha : \mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Q}_\ell$ as above. Let $\mathcal{G}$ denote the local system on S , defined by

(3.2.1)

$$\mathcal{V}_2 \otimes (\mathrm{MC}_\chi(\mathcal{V}_1 \otimes \mathrm{MC}_\chi(\mathcal{V}_2 \otimes \mathrm{MC}_\chi(\mathcal{V}_1 \otimes \mathrm{MC}_\chi(\mathcal{V}_2 \otimes \mathrm{MC}_\chi(\mathcal{V}_1 \otimes \mathrm{MC}_\chi(\mathcal{V}_0))))))).$$

In [6], the following is shown:

3.2.1 Proposition. *The monodromy tuple of $\mathcal{G}$ is of the form $(\rho_{\mathcal{G}}(\delta_1), \rho_{\mathcal{G}}(\delta_2)) \in \mathrm{GL}_7(\mathbb{Z})$, where $(\rho_{\mathcal{G}}(\delta_1), \rho_{\mathcal{G}}(\delta_2))$ is as follows:*

$$\left(\begin{pmatrix} 1 & 0 & 0 & 2 & 2 & 0 & 0 \\ 0 & 1 & 0 & -2 & 0 & 2 & 0 \\ 0 & 0 & 1 & 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 2 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 & 4 & 4 & 1 \end{pmatrix} \right).$$

For any prime $\ell > 3$, the ℓ -adic closure of the monodromy group coincides with the group $G_2(\mathbb{Z}_\ell)$. Especially, the Zariski closure of the monodromy group in $\mathrm{GL}_7(\mathbb{C})$ coincides with the group $G_2(\mathbb{C})$.

Now, consider the following Fuchsian rank one systems F_i , $i = 0, 1, 2$:

$$F_0 : Y' = \left(-\frac{1}{2t} - \frac{1}{2(t-1)} \right) Y, \quad F_1 : Y' = \frac{1}{2(t-1)} Y = \left(\frac{0}{2t} + \frac{1}{2(t-1)} \right) Y$$

and

$$F_2 : Y' = \frac{1}{2t} Y = \left(\frac{1}{2t} + \frac{0}{2(t-1)} \right) Y.$$

The local systems of holomorphic solutions of F_i , $i = 0, 1, 2$, on $S = \mathbb{A}^1 \setminus \{0, 1\}$ can easily be seen to be isomorphic to the above local systems $\mathcal{V}_i$, $i = 0, 1, 2$.

The tensor product of two Fuchsian systems associated to $(A_1, \dots, A_r)$ and $(B_1, \dots, B_r)$ is given by the Fuchsian system associated to the tuple

$$(A_1 \otimes 1 + 1 \otimes B_1, \dots, A_r \otimes 1 + 1 \otimes B_r).$$

The dual of a system $F : Y' = CY$ of linear differential equations is given by $F^* : Y' = -C^{\text{tr}}Y$, where C^{tr} denotes the transpose of the matrix C .

Let G be the Fuchsian system which is given by the following sequence of middle convolutions and tensor products:

(3.2.2)

$$F_2^* \otimes (\text{mc}_{-\frac{1}{2}}(F_1^* \otimes \text{mc}_{\frac{1}{2}}(F_2^* \otimes \text{mc}_{-\frac{1}{2}}(F_1 \otimes \text{mc}_{\frac{1}{2}}(F_2 \otimes \text{mc}_{-\frac{1}{2}}(F_1^* \otimes \text{mc}_{\frac{1}{2}}(F_0))))))))).$$

Recall from [14], that any system of linear differential equations

$$F : Y' = AY, \quad A \in \mathbb{C}(t)^{n \times n},$$

has attached an algebraic group G_F to it, called the *differential Galois group* of F . Moreover, if the singularities of F are regular, then G_F can be viewed as the Zariski closure of the monodromy of F inside the group $\text{GL}_n(\mathbb{C})$. With this, we obtain the following result:

3.2.2 Theorem. *The Fuchsian system G is given by*

$$Y' = \left(\frac{1}{2t} \cdot \begin{pmatrix} -2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix} + \frac{1}{2(t-1)} \cdot \begin{pmatrix} 0 & 0 & 0 & 0 & -2 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 & -2 \end{pmatrix} \right) Y.$$

The differential Galois group of G is isomorphic to $G_2(\mathbb{C})$. Moreover, the Fuchsian system G is globally nilpotent.

Proof: The form of the matrices follows from the explicit construction of mc_μ . By the Riemann-Hilbert correspondence for mc_μ (Thm. 1.2.3), the monodromy tuple of G is given by the matrices $(\rho_{\mathcal{G}}(\delta_1), \rho_{\mathcal{G}}(\delta_2))$ above. Thus the first claim follows from Prop. 3.2.1 and the fact that the differential Galois group of a Fuchsian system is the Zariski closure of the monodromy (since all singularities are regular), see [14]. The last claim follows from Thm. 2.0.6. $\square$

3.2.3 Remark. The above Fuchsian system can be viewed to be the de Rham realization of a certain mixed motive M of dimension 7, which appears in [6]. The motive M stands in close connection to a question of Serre on the existence of motives with exceptional Galois group G_2 , see [11], [6].

References

- [1] Y. André. *G-Functions and Geometry*. Aspects of Mathematics 13. Vieweg, 1989.
- [2] P. Deligne. *Equations Différentielles à Points Singuliers Réguliers*. Lecture Notes in Mathematics 163. Springer-Verlag, 1970.
- [3] M. Dettweiler. Galois realizations of classical groups and the middle convolution. *Habilitation Thesis, Heidelberg*, 2005.
- [4] M. Dettweiler and S. Reiter. An algorithm of Katz and its application to the inverse Galois problem. *J. Symb. Comput.*, 30: 761–798, 2000.
- [5] M. Dettweiler and S. Reiter. On the middle convolution. *Preprint math.AG/0305311*, 2003.
- [6] M. Dettweiler and S. Reiter. Rigid local systems and relative motives with Galois group G_2 . *Preprint*, 2006.
- [7] C. Doran. Algebraic and geometric isomonodromic deformations. *J. Differential Geom.*, 59: 33–85, 2001.
- [8] N. Katz. Nilpotent connections and the monodromy theorem: applications of a result of Turrittin. *Publ. Math. IHES*, 39: 175–232, 1970.
- [9] N.M. Katz. *Rigid Local Systems*. Annals of Mathematics Studies 139. Princeton University Press, 1996.
- [10] D. Krammer. An example of an arithmetic Fuchsian group. *J. reine angew. Math.*, 473: 69–85, 1996.

- [11] J.-P. Serre. Propriétés conjecturales des groupes de Galois motiviques et des représentations ℓ -adiques. In U. Jannsen et al., editor, *Motives I*, number 55 in Proceedings of Symposia in Pure Mathematics, pages 377–400, 1994.
- [12] M. van der Put. Reduction modulo p of differential equations. *Indag. Math.*, 7: 367–387, 1996.
- [13] M. van der Put. Grothendieck’s conjecture for the Risch equation $y' = ay + b$. *Indag. Math.*, 12: 113–124, 2001.
- [14] M. van der Put and M. Singer. *Galois theory of linear differential equations*, volume 328 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 2003.

Michael Dettweiler
 IWR, Universität Heidelberg,
 INF 368
 69121 Heidelberg, Deutschland
 e-mail: michael.dettweiler@iwr.uni-heidelberg.de

Stefan Reiter
 Technische Universität Darmstadt
 Fachbereich Mathematik AG 2
 Schloßgartenstr. 7
 64289 Darmstadt, Deutschland
 email: reiter@mathematik.tu-darmstadt.de